

Some Theorems on Bounded Analytic Functions Author(s): Walter Rudin Source: Transactions of the American Mathematical Society, Vol. 78, No. 2 (Mar., 1955), pp. 333-342 Published by: American Mathematical Society Stable URL: <u>http://www.jstor.org/stable/1993067</u> Accessed: 30/11/2009 01:57

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/action/showPublisher?publisherCode=ams.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to Transactions of the American Mathematical Society.

SOME THEOREMS ON BOUNDED ANALYTIC FUNCTIONS

BY

WALTER RUDIN

1. Introduction. In a paper by Bers [3] the following unpublished theorem of Chevalley and Kakutani is mentioned in a footnote:

THEOREM A. If to each boundary point W of a domain B there exists a bounded analytic function defined in B and possessing at W a singularity, then B is determined (modulo a conformal transformation) by the ring of all bounded analytic functions on B.

The purpose of this paper is twofold. A proof of Theorem A is given which, according to Professor Kakutani, is considerably simpler than the original one (Theorem 9; a slight extension is stated in Theorem 12). In particular, no use is made of the theory of normed rings and of the topology of the maximal ideal space; our methods are more function-theoretic. Secondly, we investigate those properties of the boundary of a domain which are connected with the existence of singularities of bounded analytic functions. This leads to the consideration of two classes of boundary points (removable and essential) and of their relation to Painlevé null-sets (Theorem 5). A theorem on cluster sets is established (Theorem 14), and it is shown (Theorem 15) that the domains considered by Chevalley and Kakutani (we shall call these domains maximal) are precisely those which are domains of existence of single-valued bounded analytic functions, i.e., those whose boundaries are natural boundaries of some such function. Theorem 11 shows that every domain D is contained in a unique smallest maximal domain D^* .

2. Definitions. Let B(D) denote the set of all single-valued bounded analytic functions on the domain D (by a domain we mean a connected open subset of the Riemann sphere S; unless the contrary is stated, we shall always assume that the boundary of D is bounded; this involves no loss of generality, since all our results are invariant under conformal one-to-one mappings of Sonto S).

Let $f \in B(D)$. If D_1 is a domain such that $D_1 \cap D$ is not vacuous, and if there is a function $f_1 \in B(D_1)$ such that $f(z) = f_1(z)$ for $z \in D_1 \cap D$, we say that f can be *extended* to D_1 .

A boundary point x of D is said to be *removable* if for every $f \in B(D)$ there exists a neighborhood V of x such that f can be extended to V. By an *essential* boundary point of D we mean one that is not removable. If every boundary

Presented to the Society, December 29, 1953, under the titles Natural boundaries of bounded analytic functions and Rings of bounded analytic functions; received by the editors January 7, 1954.

point of D is essential, we say that D is maximal.

The closed disc with center at x and radius r will be denoted by J(x, r).

3. Painlevé null-sets. Let K be the complement of a domain D. If B(D) consists of the constant functions alone, then K is called a Painlevé null-set [2, p. 107] or simply a null-set. It is known that K is a null-set if the linear measure of K is zero [1, p. 2]; if K is a subset of an analytic arc, this sufficient condition is also necessary [2, p. 122]; in the general case, no characterization of this sort has yet been found.

It is trivial that null-sets are totally disconnected, and it is easily shown [2, p. 108] that K is a null-set if and only if every $f \in B(\Delta - K)$ can be extended to Δ , where Δ is any domain containing K. This remark leads to a simple proof of the following property, which we state as a lemma:

4. LEMMA. Let Δ be a domain, and let K be a closed set. Suppose that for every $x \in \Delta \cap K$ there exists r > 0 (r depending on x) such that $J(x, r) \subset \Delta$ and $J(x, r) \cap K$ is a Painlevé null-set. Then every $f \in B(\Delta - K)$ can be extended to Δ .

Next we show that if x is a removable boundary point of D, there exists a domain D_1 containing x and D, *independent* of f, such that every $f \in B(D)$ can be extended to D_1 :

5. THEOREM. Let K be the complement of the domain D. A point $x \in K$ is a removable boundary point of D if and only if the set $K \cap J(x, r)$ is a Painlevé null-set for some r > 0.

Proof. The sufficiency of the condition is evident. To prove the necessity, suppose $K \cap J(x, r)$ is not a null-set for any r > 0. Let V be a neighborhood of x, and put $\Delta = V - \{x\}$. If for every $y \in \Delta$ there exists an r > 0 such that $K \cap J(y, r)$ is a null-set, then every $f \in B(\Delta - K)$ can be extended to Δ , by Lemma 4, and hence to V, so that $K \cap J(x, r)$ is a null-set provided $J(x, r) \subset V$.

This contradiction shows that there exists a sequence of disjoint discs $J(x_n, r_n) \subset \Delta$ $(n = 1, 2, 3, \cdots)$ such that $x_n \to x$ and such that $K_n = K \cap J(x_n, r_n)$ is not a null-set. The complement D_n of K_n contains D and is connected (being the union of two connected sets). Choose nonconstant functions $f_n \in B(D_n)$ such that $|f_n(z)| < 2^{-n}$ in D_n , and define $f(z) = \sum f_n(z)$.

Then $f \in B(D)$. Since $f = (f - f_n) + f_n$ and $f - f_n$ can be extended to a neighborhood of K_n , whereas this is not true of f_n , we see that f cannot be extended to any neighborhood of x. Hence x is an essential boundary point of D, and the theorem follows.

The following two lemmas will be needed in the proof of Theorem 9.

6. LEMMA. Suppose Δ is an open set which is dense in a domain D. If ϕ is analytic in D (with possible poles) and schlicht in Δ , then ϕ is schlicht in D.

Proof. Suppose there are two distinct points z_1 , z_2 in D such that $\phi(z_1)$

 $=\phi(z_2)$, and let V_1 , V_2 be disjoint neighborhoods of z_1 , z_2 , such that $V_1 \subset D$, $V_2 \subset D$.

Since $\phi(V_i \cap \Delta)$ is an open set, dense in $\phi(V_i)$ (i=1, 2), we see that $\phi(V_1 \cap \Delta)$ and $\phi(V_2 \cap \Delta)$ have a point in common; this point has two distinct inverse images in Δ , a contradiction.

7. LEMMA. Let D_1 be a domain. If D_1 is maximal and conformally equivalent to D_2 , then D_2 is also a maximal domain.

Proof. Suppose ϕ maps D_2 conformally onto D_1 . We assume, without loss of generality, that ϕ is analytic, rather than conjugate analytic. If D_2 has a removable boundary point x, there exists a neighborhood V of x such that every $f \in B(D_2)$ can be extended to $D'_2 = V \cup D_2$, by Theorem 5 and Lemma 4. D_2 is clearly dense in D'_2 , and since ϕ is bounded except in the neighborhood of the point $z_0 \in D$ such that $\phi(z_0) = \infty$ (if D_1 contains the point at infinity), ϕ can be extended to D'_2 . By Lemma 6, ϕ is schlicht in D'_2 , so that the domain $D'_1 = \phi(D'_2)$ contains D_1 properly.

Choose $f \in B(D_1)$, and set $g(w) = f(\phi(w))$ for $w \in D_2$. Since $g \in B(D_2)$, g can be extended to D'_2 , and the formula: $f(z) = g(\phi^{-1}(z))$ extends f to D'_1 , contradicting the maximality of D.

Hence D_2 has no removable boundary point.

8. Before turning to the proof of the theorem of Chevalley and Kakutani, we insert the following remark: We shall consider two domains, D_1 and D_2 , and an isomorphic mapping ψ of the ring $B(D_1)$ onto the ring $B(D_2)$, i.e., a one-to-one mapping which preserves sums and products; it is clear that $\psi(1) = 1$. Since the imaginary unit *i* is a primitive fourth root of unity, we must either have $\psi(i) = i$ or $\psi(i) = -i$. In the former case, we call ψ a *direct* isomorphism. If $\psi(i) = -i$, conjugates must be introduced into (1) below in an obvious manner, and the induced correspondence between the points of D_1 and D_2 , while still conformal, reverses orientation.

9. THEOREM. Let D_1 and D_2 be maximal domains which are proper subsets of the Riemann sphere, and let ψ be a direct isomorphism mapping $B(D_1)$ onto $B(D_2)$. Then there exists a one-to-one conformal mapping ϕ of D_2 onto D_1 such that

(1)
$$f(\phi(w)) = f^*(w) \qquad (w \in D_2),$$

where $f^* = \psi(f)$.

Since the converse of the theorem is obvious, we can summarize the situation by saying that D_1 and D_2 are conformally equivalent if and only if $B(D_1)$ and $B(D_2)$ are algebraically isomorphic.

Proof. We assume, without loss of generality, that D_1 and D_2 are in the finite plane.

WALTER RUDIN

Using the notation $f^* = \psi(f)$, we readily see that $r^* = r$ for any rational number r (i.e., for any constant function whose range is the number r). Having assumed that $i^* = i$, we see that $r^* = r$ for any complex rational r (by a complex rational we mean a complex number with rational real and imaginary parts).

Let Q(f) denote the closure of the range of f. For any $f \in B(D_1)$ and any complex rational r, the following four statements are clearly equivalent, since

$$(f - r)^* = f^* - r^* = f^* - r$$

and ψ preserves reciprocals:

(a) $r \in Q(f)$.

(b) f-r has no reciprocal in $B(D_1)$.

(c) $f^* - r$ has no reciprocal in $B(D_2)$.

(d) $r \in O(f^*)$.

Thus Q(f) and $Q(f^*)$ contain the same complex rationals.

It follows that for any constant c, c^* is also a constant. Furthermore, if f is not constant, then the same is true of f^* ; since Q(f) and $Q(f^*)$ are closures of open sets, we conclude that

(2)
$$Q(f) = Q(f^*) \qquad (f \in B(D_1), f \neq \text{const.}).$$

Now let c be any constant, choose a nonconstant $f \in B(D_1)$ such that $0 \in Q(f)$, and put

$$f_n(z) = c + f(z)/n$$
 $(z \in D_1, n = 1, 2, 3, \cdots).$

Then

$$f_n^*(w) = c^* + f^*(w)/n$$
 $(w \in D_2, n = 1, 2, 3, \cdots),$

and $c \in Q(f_n)$, $c^* \in Q(f_n^*)$. By (2) we have

$$c = \bigcap_{n=1}^{\infty} Q(f_n) = \bigcap_{n=1}^{\infty} Q(f_n^*) = c^*,$$

so that $c^* = c$ for every constant c.

Now, if D_1 is unbounded, the identity function I(z) = z is not a member of $B(D_1)$. This difficulty cannot, in general, be eliminated by mapping D_1 conformally onto a bounded domain: Ahlfors and Beurling [2] have shown that there are domains D on which there exist no bounded schlicht functions, although B(D) contains nonconstant functions; such a domain D is not conformally equivalent to any bounded domain.

However, let $F(D_j)$ be the quotient field of $B(D_j)$ (j=1, 2), and extend ψ in the natural manner to an isomorphic mapping of $F(D_1)$ onto $F(D_2)$. We shall see that $I \in F(D_1)$ and that the function $\phi = \psi(I) \in F(D_2)$ is the desired mapping function for which (1) holds.

For any nonconstant $f \in B(D_1)$ and any point $a \in D_1$, put

$$g(z) = (f(z) - f(a))/(z - a)$$

so that

$$I(z) = z = a + (f(z) - f(a))/g(z).$$

These formulas show that $g \in B(D_1)$ and hence that $I \in F(D_1)$.

Now set $\phi = \psi(I)$, $\phi(D_2) = D_3$, $D = D_1 \cap D_3$. For any constant *c*, the following four statements are clearly equivalent:

(a) $c \in Q(I)$.

(b) The reciprocal of I-c is not in $B(D_1)$.

(c) The reciprocal of $\phi - c$ is not in $B(D_2)$.

(d) $c \in Q(\phi)$.

Thus D_1 and D_3 have the same closure; it follows that D is dense in D_1 and in D_3 .

We have to show that ϕ is schlicht, and that $D_1 = D_3$.

Choose any point $a \in D$ and let M_a be the set of all functions $f \in B(D_1)$ such that f(a) = 0. M_a is evidently an ideal. If $g \in B(D_1)$ and $g(a) \neq 0$, then the identity

$$1 = \frac{g(a) - g(z)}{g(a)} + \frac{g(z)}{g(a)}$$

shows that every ideal containing M_a and g contains the constant 1 and hence the whole ring $B(D_1)$. Thus M_a is a maximal ideal of $B(D_1)$, which ψ maps onto a corresponding maximal ideal M_a^* of $B(D_2)$.

For every $f \in M_a$, $f/(I-a) \in B(D_1)$, so that $f^*/(\phi-a) \in B(D_2)$ for every $f^* \in M_a^*$. It follows that $f^*(w) = 0$ whenever $\phi(w) = a$, for every $f^* \in M_a^*$. If there were more than one such point $w \in D_2$, the ideal M_a^* would not be maximal. We conclude: for every $a \in D$, there exists precisely one point $b \in D_2$ such that $\phi(b) = a$.

If Δ is the set of all $b \in D_2$ such that $\phi(b) \in D$, the fact that D is dense in D_3 implies that Δ is dense in D_2 , since the analytic function ϕ is an open mapping. Lemma 6 now shows that ϕ is schlicht.

Thus D_3 is conformally equivalent to D_2 ; by Lemma 7, D_3 is maximal.

To complete the proof, choose $f \in B(D_1)$, $a \in D$, $b \in D_2$ such that $\phi(b) = a$. Then $f-f(a) \in M_a$, so that $f^*-f(a) \in M_a^*$. But M_a^* consists of those functions of $B(D_2)$ which vanish at b. Thus $f^*(b) = f(a)$, or

(3)
$$f(z) = f^*(\phi^{-1}(z))$$
 $(f \in B(D_1), z \in D).$

Since ϕ maps D_2 conformally onto D_3 , there is a function $g \in B(D_3)$ such that

(4)
$$g(z) = f^*(\phi^{-1}(z))$$
 $(z \in D_s).$

The functions f and g coincide in D; thus every $f \in B(D_1)$ can be extended to D_3 .

On the other hand, for every $g \in B(D_3)$ there is an $f^* \in B(D_2)$ for which (4) holds; defining $f = \psi^{-1}(f^*)$, we obtain a member of $B(D_1)$ for which (3) holds; thus every $g = B(D_3)$ can be extended to D_1 .

Since D_1 and D_3 are maximal domains, we conclude that $D_1=D_3$. The theorem follows.

10. REMARK. Let us apply Theorem 9 to the case $D_1 = D_2 = D$, say. Let ψ_1, ψ_2 be two direct automorphisms of B(D), and let ϕ_1, ϕ_2 be the corresponding conformal transformations of D onto itself, in accordance with formulas (1). Then it is easily seen that the automorphism $\psi_2\psi_1$ corresponds to the mapping $\phi_1\phi_2$, i.e., the mapping taking z into $\phi_1(\phi_2(z))$. Replacing the automorphisms ψ by their inverses, the following is evident:

For every maximal domain D, there is a natural isomorphism between the group of all direct automorphisms of the ring B(D) and the group of all directly conformal one-to-one mappings of D onto D.

We now return to our investigation of the properties of essential boundary points.

11. THEOREM. If K^* is the set of all essential boundary points of a domain D, then K^* is perfect (if not vacuous). If D^* is that component of the complement of K^* which contains D, then D is dense in D^* , and D^* is the smallest maximal domain which contains D.

Proof. Theorem 5 shows that D^* may be obtained in the following manner: cover every removable boundary point x of D with an open disc U_x which does not intersect K^* ; D^* is the union of D and these discs U_x . Since null-sets are totally disconnected, D is dense in D^* .

By Lemma 4, every $f \in B(D)$ can be extended to D^* . Thus there is no smaller maximal domain containing D. To show that D^* is maximal, let x be a boundary point of D^* . Then x is an essential boundary point of D, there is some $f \in B(D)$ which cannot be extended to any neighborhood of x, and the same is of course true of the extension of f to D^* . Thus x is an essential boundary point of D^* .

Thus D^* is maximal, and since isolated boundary points are removable, K^* is perfect (or vacuous).

As a consequence of Theorem 11, Theorem 9 may now be restated as follows:

12. THEOREM. Let D_1^* and D_2^* be the smallest maximal domains containing the domains D_1 and D_2 . If $B(D_1)$ and $B(D_2)$ are algebraically isomorphic, then D_1^* and D_2^* are conformally equivalent.

13. DEFINITION. Let x be a boundary point of a domain D, and let f be analytic in D. Put $D_n = D \cap J(x, 1/n)$, and let E_n be the closure of $f(D_n)$ (the range of the restriction of f to D_n). The set

338

$$C(f; x) = \bigcap_{n=1}^{\infty} E_n$$

is called the *cluster set* of f at x.

It is clear that for every $w \in C(f; x)$ there is a sequence $\{z_n\}$ such that $z_n \in D$, $z_n \to x$, and $f(z_n) \to w$ as $n \to \infty$.

If x is an essential boundary point of D, then there is some $f \in B(D)$ with a singularity at x. The question as to how "bad" this singularity can be is partially answered by our next theorem; the example given at the end of this paper has its origin in the same question.

14. THEOREM. Let x be an essential boundary point of the domain D. There exists a function $f \in B(D)$ whose cluster set at x consists of the entire closed unit disc, although |f(z)| < 1 for every $z \in D$.

Proof. Since the set of essential boundary points of D is perfect, there is a sequence $\{x_n\}$ of distinct essential boundary points such that $x_n \neq x$ and $x_n \rightarrow x$ as $n \rightarrow \infty$. Choose $\{r_n\}$ such that no two of the discs $T_n = J(x_n, r_n)$ intersect and such that T_n contains no point of the interior I of K if x_n is not a limit point of I. Let $K_n = K \cap J(x_n, r_n/3)$, let G_n be the complement of K_n , and put $d_n = D \cap J(x_n, r_n/2)$. As usual, K denotes the complement of D.

We shall construct an increasing sequence $\{n_k\}$ of positive integers, and functions $f_k \in B(G_{n_k})$ $(k=1, 2, 3, \cdots)$ such that

(i) $|f_k(z)| < 1$ for $z \in G_{n_k}, f_k(x) = 0$;

(ii) for some sequence $\{z_i\}$ of points in $d_{n_k}, f_k(z_i) \rightarrow 1$ as $i \rightarrow \infty$;

(iii) $|f_k(z)| < 2^{-k-1}$ for $z \in G_{n_k} - T_{n_k}$;

(iv) $\sum_{i=1}^{k} |f_i(z)| < 2^{-k-2}$ for $z \in T_{n_{k+1}}$.

If x_n is not a limit point of I, choose a nonconstant $u_n \in B(G_n)$ such that $\sup |u_n(z)| = 1$ for $z \in G_n$, hence also for $z \in d_n$, since d_n is dense in G_n .

If x_n is a limit point of *I*, choose $y_n \in I$ such that $|x_n - y_n| < r_n/6$, let p_n be the distance from y_n to *D*, and set $u_n(z) = p_n/(z-y_n)$.

In either case, let us multiply u_n by a suitable constant of modulus 1 so as to make $u_n(x)$ real, and let us define

$$v_n(z) = \frac{u_n(z) - u_n(x)}{1 - u_n(x)u_n(z)}$$

Then $v_n \in B(G_n)$, $|v_n(z)| < 1$ in G_n , and $v_n(x) = 0$; multiplying v_n by a suitable constant of modulus 1, we may further arrange it so that $v_n(z_i) \rightarrow 1$ for some sequence $\{z_i\}$ in d_n .

Now take $n_1=1$, $g_1=v_1$. Having picked n_1, \dots, n_k and g_1, \dots, g_k , take $n_{k+1} > n_k$ such that (iv) holds with g_i in place of f_i (this is possible since $g_i(x) = 0$ and $x_i \rightarrow x$), and put $g_{k+1}=v_{n_{k+1}}$.

Finally, let $f_k = (g_k)^{m_k}$, where m_k is a positive integer chosen large enough for (iii) to hold. Then all of the conditions (i) to (iv) are satisfied.

[March

Next, let *E* be a countable dense subset of the open unit disc, and let $\{w_n\}$ be a sequence of complex numbers which contains every member of *E* infinitely many times, with the additional requirement that $|w_n| < 1 - 2^{-n}$.

The function

$$f(z) = \sum_{k=1}^{\infty} w_k f_k(z) \qquad (z \in D)$$

has the desired properties. By (iii) and (iv) we have, for $z \in D \cap T_{n_m}$,

$$| f(z) | < \sum_{1}^{m-1} | f_k(z) | + | w_m | + \sum_{m+1}^{\infty} | f_k(z) |$$

 $< 2^{-m-1} + 1 - 2^{-m} + \sum_{m+1}^{\infty} 2^{-k-1} = 1.$

If $z \in D - T_{n_m}$ for every *m*, then $|f(z)| < \sum 2^{-k-1} = 1/2$. Thus |f(z)| < 1 for every $z \in D$.

Finally, choose $z \in d_{n_m}$ such that $|f_m(z) - 1| < 2^{-m}$. Then

$$|f(z) - w_m| < \sum_{1}^{m-1} |f_k(z)| + |w_m| \cdot |f_m(z) - 1| + \sum_{m+1}^{\infty} |f_k(z)| < 2^{-m+1}.$$

Thus every member of E is in C(f;x); since C(f;x) is evidently closed, the theorem follows.

15. THEOREM. If D is a maximal domain, there exists a function $f \in B(D)$ which has the boundary of D as its natural boundary.

Proof. Suppose, without loss of generality, that D contains the point at infinity. Let $\{x_n\}$ be a countable dense subset of the boundary of D. The preceding theorem, applied to each x_n , shows that there exist functions $f_n \in B(D)$ and points $z_n \in D$ such that $|f_n(z)| < 1$ in D, $f_n(\infty) = 0$, $|x_n - z_n| < 1/n$, $f_n(z_n) = a_n$, where a_n is real and

$$\sum_{n=1}^{\infty} (1-a_n) < \infty.$$

Define

$$f(z) = \prod_{n=1}^{\infty} \frac{a_n - f_n(z)}{1 - a_n f_n(z)} \cdot$$

By a generalization of Schwarz's lemma, for every closed subset C of D there is a constant M < 1 such that $|f_n(z)| < M$ on C. For $z \in C$, we have

$$\left|1 - \frac{a_n - f_n(z)}{1 - a_n f_n(z)}\right| < \frac{1 + M}{1 - M} (1 - a_n),$$

340

so that the product defining f(z) converges uniformly on C. Since $f(z_n) = 0$ and every boundary point of D is a limit point of the set $\{z_n\}$, the theorem follows.

16. An example. In view of Theorem 14, one may ask the following question: given an essential boundary point x of the domain D and an arc L which lies in D, except for one of its end points, which is at x, does there exist a nonconstant function $f \in B(D)$ such that (a) f(z) does not converge as $z \rightarrow x$ along L, or, (b) such that |f(z)| tends to its maximum as $z \rightarrow x$ along L?

It can be shown that if for some x the answer to (b) is affirmative, the same is true for (a). We shall give an example in which the answers to both (a) and (b) are negative.

Let $\{a_n\}$ be a sequence of distinct points on the positive real axis, converging to zero. Choose $\{R_n\}$ such that the discs $J(a_n, R_n)$ are mutually disjoint, let $\{c_n\}$ be a sequence of positive numbers such that $c_n < 1$ and

$$\sum_{n=1}^{\infty} c_n/(1-c_n) < \epsilon,$$

where $\epsilon > 0$ is given, put $r_n = c_n R_n$, $V_n = J(a_n, r_n)$, let K consist of the origin and the union of the V_n 's, and let D be the complement of K. Since every boundary point of every V_n is an essential boundary point of D and the set of essential boundary points is closed, the origin is an essential boundary point of D.

Let $f \in B(D)$, and suppose |f(z)| < 1, $f(\infty) = 0$. At almost all points of the circumference C_n of each V_n , f(z) has well-defined nontangential boundary values, so that the path of integration in the Cauchy integral formula can be moved to C_n . If we put

$$f_n(z) = \frac{1}{2\pi i} \int_{C_n} \frac{f(t)}{t-z} dt \qquad (z \notin V_n),$$

then it is easily verified that

$$f(z) = \sum_{n=1}^{\infty} f_n(z),$$

the series converging uniformly in every closed subset of D. Moreover, if $z \notin G_n$, where G_n is the interior of $J(a_n, R_n)$, we have

$$\left| f_n(z) \right| \leq \frac{r_n}{R_n - r_n} = \frac{c_n}{1 - c_n}$$

If $G = \bigcup G_n$, it follows that the series defining f(z) converges absolutely and uniformly on the complement H of G. Thus f(z) is continuous on H, which includes the origin, and $|f(z)| \leq \epsilon$ on H.

WALTER RUDIN

If L is now any arc in H, with one end point at the origin, then $\lim f(z)$ exists as $z \to 0$ along L, and does not exceed ϵ in absolute value. This is so even if the cluster set of f at 0 consists of the entire unit disc.

References

1. L. V. Ahlfors, Bounded analytic functions, Duke Math. J. vol. 14 (1947) pp. 1-11.

2. L. V. Ahlfors and Arne Beurling, Conformal invariants and function-theoretic null-sets, Acta Math. vol. 83 (1950) pp. 101-129.

3. Lipman Bers, On rings of analytic functions, Bull. Amer. Math. Soc. vol. 54 (1948) pp. 311-315.

UNIVERSITY OF ROCHESTER, Rochester, N. Y.